

01.12.2016. , 5940/8,

:

”

“

”

“

”

1.

2.

3.

) \_\_\_\_\_

149 a

: , , , ,

, . 25

,17 86 .

, ,

.

,

.

,

.

.

.

,

.

.

,

.

,

,

:

,

,

,

.

.

.

.

,

.

:

,

: )

)

)

)

-

”

“

78

103

,

(KneeSocietyClinicalRatingSystem

WesternOntarioandMcMasterUniversitiesArthritisIndex).

(KneeSocietyRoentgenographicEvaluationSystem).

3,6

12

) \_\_\_\_\_

(Knee Society Score– *KSS*),  
 18±11,4 83±3,5,  
 (Functional Knee Society Score –*FKSS*) a 21±9,9 50±5,9,  
 (Western Ontario and McMaster Universities Arthritis Indeks-  
*WOMAC*) 23±6,4 69±4,6. *KSS*  
 26±9,0 82±2,9, *FKSS* 27±9,7 50±7,4,  
*WOMAC* 27±6,3 a 69±5,0.

je

0,9%,

96,3%,

94,2%,

t (0.9%), a

3

(2.5%)

(p=0.407).

80

97,2%, a

95,8%.

(BAC),

>20, 12,9%

<20

>20, 21,2%

<20,

>80, 1,1%

<80, 16,7%

<80, 9,5%

>80

100%.

KSS FKSS,  
WOMAC  
( $r < 0,5$ ).

)

---

8  
(J.K.Lee,  
2012).

Almedia  
(2008), Ranawat (1989), Dalury (1995),  
Stundner (2014), 4,5%  
3,5%  
0,9%, 36  
Rodriguez  
(1996) e  
4,1%,  
7

(Bongartz 2008, Schrama 2010),

, , 1,26%  
0,84%, Bheeshma Ravi (2014).

(Denglu, 2013).

, ,  
11 ,  
:  
94,2%. o  
10  
: 97,34% (Scuderi, 1989), 97% (Schai, 1999), 93%  
(Hanyu, 1997), 96% (Himanen, 2005).

, ,  
(Yoshishara 2016, Goodman 2016).

, 20 ,  
71,3% 72,7%

( 84,7 77,7). (Noble 2006, Bourne 2010, Dunbar 2013).

WOMAC . , KSS FKSS

)  
*Blagojevi Z, Gluš evi B, or evi B. Total knee arthroplasty in patients with rheumatoid arthritis-midterm results  
Vojnosanitetski pregled, 2016 OnLine-First (00):154-154  
DOI:10.2298/VSP150901154G*

) ( )  
” “ , ao



,  
,  
.  
,2016.  
:  
.  
.  
.  
.